

December, 1996
 LPTENS-96/64
 UT-Komaba-96/24

FANCY AND FACTS IN THE $(d - 2)$ EXPANSION OF NON-LINEAR SIGMA MODELS

E. Brézin^{a)} and S. Hikami^{b)}

^{a)} Laboratoire de Physique Théorique, Ecole Normale Supérieure
 24 rue Lhomond 75231, Paris Cedex 05, France¹

^{b)} Department of Pure and Applied Sciences, University of Tokyo
 Meguro-ku, Komaba, Tokyo 153, Japan

Abstract

We review the existing results on the scaling dimensions of operators with more than two derivatives in the non-linear sigma models. We argue that the speculations on the relevance of these operators, and correspondingly on the breakdown of the $(d - 2)$ expansion for the classical Heisenberg model, or for the one-parameter scaling theory of localization, are based on a dubious mathematical analysis.

The low temperature expansion of the N -vector model is given by a model of interacting Goldstone bosons, known as the non-linear sigma model [1, 2]. The existence of a zero-temperature fixed point in dimension two has led to an expansion of the critical properties in powers of $(d - 2)$ which supplements the familiar expansion in powers of $(4 - d)$ based on the linear Landau-Ginzburg-Wilson model [3]; linear and non-linear refer here to the representation of the $O(N)$ symmetry by the order parameter.

These non-linear sigma models were generalized to situations in which the order parameter is a mapping of a two-dimensional lattice onto a target manifold (a sphere in the case of the $O(N)$ model). These generalizations are the basis of the conformal field theory approach to string theory, and they

¹ Unité propre du Centre National de la Recherche Scientifique, associée à l'Ecole Normale Supérieure et à l'Université de Paris-Sud

have been also applied to various problems in condensed matter physics, in particular to Anderson localization [4], for which the manifold is the homogeneous space $O(N, N)/O(N) \times O(N)$ [5, 6, 7], in the replica limit in which N goes to zero. This had led to an expansion of the metal-insulator transition in dimension three in powers of $\epsilon = d - 2$, with $\epsilon = 1$ at the end. The numerical success of this expansion has been unfortunately in no way comparable to that of the Wilson-Fisher expansion in powers of $(4 - d)$. However for the localization problem, there is no upper critical dimension from which an expansion similar to the $(4 - d)$ would allow us to ignore the poor numerical accuracy of the $(d - 2)$ expansion. There is no other analytic approach to the problem.

In view of these difficulties a growing number of articles, starting with the work of Kravtsov, Lerner and Yudson [8], extended later to the more familiar $O(N)$ model by Wegner [9], pointed out to the possible appearance of new relevant operators, which would of course drastically change the physics and the results of the model.

In this note we would like to put the calculations which led to the alleged breakdown of the $(d - 2)$ expansion, in the right perspective. Not that we have any significant additional calculation to report, but we would like the reader to judge the seriousness of the problem by presenting a summary of the situation as it is known at present.

We shall center our discussion on the $O(N)$ model whose physics is under much better control, and for which the $4 - d$ and $1/N$ expansion, plus a wealth of numerical experiments leave little doubts on its properties. Let us start by reminding the reader that the model is defined by a lattice of unit N -vectors with nearest neighbour interactions of the form $\vec{S}(x) \cdot \vec{S}(x + ae_\alpha)$, in which e_α is a unit vector in the direction α . In the long distance limit we can replace $\vec{S}(x)$ by a continuum field and the interaction by $\vec{S} \cdot \nabla^2 \vec{S}$, with the constraint $\vec{S}^2 = 1$; (the term involving one derivative of $\vec{S}$ cancels with the $x - ae_\alpha$ neighbour). The model is then defined by a straight low temperature expansion over some ordered state. However this expansion is plagued with short-distance divergences, as in any continuum field theory, and these divergences have to be regulated. This is slightly tricky in this problem; a straight ultra-violet cut-off would not work since it breaks the $O(N)$ invariance of the model. There are two schemes known for regulating these divergences without breaking the rotational invariance, the dimensional regularization or a

lattice, which means simply returning to the low temperature expansion of the original classical Heisenberg model. Of course these two regularizations lead at the end to identical results in the critical long-distance limit [2]. In other words one should be clear at this stage: if one does find a problem with this theory it is not simply a problem of some abstract non-linear field theory, but of the low temperature expansion of the Heisenberg model itself.

The difficulties which have bothered a number of workers concern the scaling behaviour of operators with more derivatives than two. Indeed it is clear from the very definition of the model that one could have operators such as $\vec{S} \cdot (\nabla^2)^2 \vec{S}$, or more complicated ones which would involve derivatives and four spins for instance, as the ones which would be generated by the interaction of four spins around a plaquette. In any way we know from Wilson theory [10] that we should allow in the Hamiltonian for any possible $O(N)$ invariant operator and find the fixed point in a space of an infinite number of coupling constants. Note however that we know a priori that there is one and only one relevant operator (in the absence of a symmetry-breaking operator such as a magnetic field); indeed it is sufficient, in order to reach the critical point to vary a single parameter, namely the temperature. If we had more relevant operators, we would have more "knobs" to adjust before we could reach criticality. Numerical experiments, expansions such as the $1/N$, have never found anything of the sort: it is sufficient to be at T_c to obtain an infinite correlation length.

In two dimensions, simple dimensional counting of operators involving $2s$ derivatives of the order parameter (an even number is needed to make a scalar), gives their scaling dimension

$$y_s = 2 - 2s \tag{1}$$

which leads to conclude that operators with more than two derivatives are irrelevant. There is no operator for $s = 0$ since $\vec{S}^2 = 1$; therefore we are left with a single relevant operator, namely $\vec{S} \cdot \nabla^2 \vec{S}$, as expected.

In dimension d two things occur : (i) the canonical dimension is changed to $(d - 2s)$ (ii) scaling anomalies appear, calculable from the renormalization group theory by expanding in powers of $(d - 2)$. Wegner's result [9], based on a one-loop calculation, gave

$$y_s = d - 2s + \frac{\epsilon s(s - 1)}{N - 2} + O(\epsilon^2) \tag{2}$$

and a recent two-loop calculation by Castilla and Chakravarty [11] gave for large s

$$y_s = d - 2s + \frac{\epsilon s(s-1)}{N-2} + \epsilon^2 \left[\frac{2}{3} \frac{s^3}{(N-2)^2} + O(s^2) \right] + O(\epsilon^3) \quad (3)$$

(their calculation is done for finite s as well but the result is only quoted for large s). Finally let us quote the last available information which is the $1/N$ expansion of Vasil'ev and Stepanenko [12] which gives

$$y_s = d - 2s + \frac{4}{N} \frac{(d-1)\Gamma(d-2)[s(s-1)d(d-3) - 2s(d-2)]}{(4-d)\Gamma(2-\frac{d}{2})\Gamma(\frac{d}{2}-1)\Gamma(\frac{d}{2}-2)\Gamma(\frac{d}{2}+1)} + O(\frac{1}{N^2}) \quad (4)$$

i.e by expanding also in powers of $\epsilon = d - 2$

$$y_s = d - 2s + \frac{\epsilon s(s-1)}{N} + \frac{\epsilon^2 s}{N} + O(\frac{\epsilon^3}{N}, \frac{1}{N^2}) \quad (5)$$

We remark here that we could determine the two loop result of y_s (3) explicitly by noting that $y_1 = 1/\nu$, where ν is a critical exponent for the correlation length, and that $y_2 = -\omega$, where ω is an exponent for the correction to the scaling. Indeed, (4) for $s = 2$ agrees with the expression of $1/N$ expansion of the exponent of the correction to the scaling by Ma [13]. From the results of $1/N$ expansion, and from (3) the two-loop term may be expressed as $(\frac{2}{3}s^3 + as^2 + (N+b)s)t_c^2$. Since we know that it is proportional to $(N-2)$, for both $s = 1$ and 2 , this fixes the two unknown constants a and b to be $a = -2$ and $b = -\frac{2}{3}$.

Thus we get

$$y_s = d - 2s + s(s-1)t_c + [\frac{2}{3}s(s-1)(s-2) + s(N-2)]t_c^2 + O(t_c^3) \quad (6)$$

where t_c is obtained from the zero of the β function [2]. Putting the expression for t_c in the ϵ expansion into (6), we have

$$\begin{aligned} y_s &= 2 + \epsilon - 2s + s(s-1)\frac{\epsilon}{N-2} \\ &+ [\frac{2}{3}s(s-1)(s-2) - s(s-1) + s(N-2)]\frac{\epsilon^2}{(N-2)^2} + O(\epsilon^3) \end{aligned} \quad (7)$$

We may recover this result by an alternative method: we consider again operators with $2s$ derivatives, but for sigma models on the target manifolds $O(N)/O(p) \times O(N-p)$. From the isomorphism between $O(5)/O(4)$ and $Sp(2)/Sp(1) \times Sp(1)$ [14], we conclude that

$$y_s = d - 2s + s(s-1)t_c + \left[\frac{2}{3}s(s-1)(s-2) + (2p(N-p) - N)s\right]t_c^2 + O(t_c^3) \quad (8)$$

using now

$$t_c = \frac{\epsilon}{N-2} - \frac{(2p(N-p) - N)\epsilon^2}{(N-2)^3} + O(\epsilon^3) \quad (9)$$

we find

$$\begin{aligned} y_s &= 2 + \epsilon - 2s + s(s-1)\frac{\epsilon}{N-2} + \left[\frac{2}{3}s(s-1)(s-2) \right. \\ &\quad \left. - \frac{2p(N-p) - N}{N-2}s(s-1) + (2p(N-p) - N)s\right]\frac{\epsilon^2}{(N-2)^2} + O(\epsilon^3) \end{aligned} \quad (10)$$

which reduces to (7) for $p = 1$.

The results of Wegner and followers led to question the possible relevance of the operators with more than two derivatives. Indeed, if in two dimensions they were irrelevant because $y_s \leq -2$ for $s > 1$, one sees that the anomaly, being quadratic in s at first order in ϵ and cubic in s at second order, could change this conclusion. Fixing d at physical dimension three, and say $N = 3$, we see that the $s = 2$ operator looks already relevant and it is a fortiori true for $s > 2$. Wegner's calculation is backed by the two-loop calculation (3) since the coefficient of ϵ^2 is also positive.

This is of course a very strange situation since it leads to the speculation that there would be an infinite number of relevant operators, whose coefficients would have to be tuned before one reaches criticality. On the basis of similar calculations for generalized sigma models, Kravtsov et al. [8], were led to conclude that the one-parameter scaling theory of Anderson et al. [15] could be invalid.

For the Heisenberg model there is no experimental or numerical indication that one could have more than one relevant operator. Therefore one would conclude that, if these higher derivatives operators were truly relevant, although there is only one fixed point at zero temperature in dimension two,

one would have to find a new fixed point in dimension greater than two which would describe the critical properties. However the $1/N$ expansion, for instance, does not show any other fixed point, although it interpolates smoothly between the $4-d$ and $d-2$ expansions. So for large N at least, this scenario with some new fixed point, would run against all the existing evidence.

Therefore let us return to the existing series (1,2); they are obtained by a standard procedure in which ϵ is a parameter which goes to zero first. The previous paradoxical conclusions are based however on fixing ϵ and letting s grow. Clearly it involves an inversion of limits. To make this point more explicit, let us focus on the existing large s information. For large s we can write the expansion (3) as

$$y_s = d - 2s[1 - \frac{1}{2}x - \frac{1}{3}x^2 + O(x^3)] \quad (11)$$

in which we have defined the parameter

$$x = \frac{\epsilon s}{N - 2} \quad (12)$$

Let us consider the function

$$f(x) = 1 - \frac{1}{2}x - \frac{1}{3}x^2 + O(x^3) \quad (13)$$

For large s , one can speculate that for higher orders in ϵ as well, the result for y_s is of the form

$$y_s = 1 - 2sf(x) \quad (14)$$

It is quite possible that for the large s the result has this form and we are trying to prove it [16]. The sign of $f(x)$, for fixed ϵ , large s , i.e. for large x , is crucial. Near $x = 0$ $f(x)$ is positive, but its asymptotic expansion at the origin, extrapolated as it is, gives a negative result. However we know how dangerous it is to study the sign of a function for large x based on the knowledge of its asymptotic expansion at small x . Even if this expansion was convergent, it would presumably have singularities somewhere, and the small x expansion would give no information on the sign of the function beyond the closest singularity. Furthermore, although there are no rigorous results, it is more than likely that the expansion in powers of x is only asymptotic

and not convergent for any x . The most naive extrapolation of $f(x)$, given the existing data, would be to replace it by a [1,1] Padé approximant, i.e. by

$$f(x) = \frac{6 - 7x}{6 - 4x} + O(x^3) \quad (15)$$

We certainly do not claim that this is in any way a valid representation of $f(x)$; however for large x , it is certainly as trustworthy as the small x -expansion (13). Of course the Padé approximation (15) is positive for large x , and the relevance of the large s -operators becomes even more doubtful.

This discussion does not constitute a proof either in any way. We wanted simply to stress how unlikely is the picture with an infinite number of relevant operators, and how weak are the mathematical assumptions on which these speculations are based.

This does not mean that the non-linear sigma models are always perfectly sound. Let us mention some real difficulties. Indeed consider an $O(N)$ -invariant Landau potential $V(\vec{\phi}^2)$ which would develop a minimum away from zero below some temperature, in other words which leads to a first order transition. In the low temperature phase of the model, there are still $(N - 1)$ Goldstone bosons, since the analysis requires simply a broken continuous symmetry, but not necessarily a second order transition. In the low temperature phase of this model, we would write a low temperature expansion, which would show a fixed point at non-zero temperature above two dimensions. Clearly it would be wrong to interpret this fixed point as a critical temperature.

The situation may even be a little more subtle. We know of cases in which the Landau potential gives a mean-field second order transition, but the fluctuations drive it to first order. This seems to happen if, for the same $O(N)$ -symmetry, the field belongs to the adjoint representation of the group, rather than to the vector one. It means that ϕ is an $N \times N$ symmetric, traceless, matrix, which transforms under the rotation ω as $\omega^T \phi \omega$. The Landau potential involves two quartic invariants, namely $(\text{Tr} \phi^2)^2$ and $\text{Tr} \phi^4$. The renormalization group at one-loop, in this space of two coupling constants, shows at first order in $4 - d$, a runaway solution to an unstable potential, indicative of a first order transition. However the analysis of the Goldstone mode of the $O(N)$ -symmetry broken down to $O(N - 1)$ depends uniquely upon the Lie algebra of these two groups, and not on the representation of the group to which belongs the order parameter. Thus, there again, we would

find a fixed point in the $d - 2$ expansion, whereas it is not at all clear that the model has a second order transition.

Therefore we do not claim that the non-linear sigma models are to be trusted as other well established field theories such as QED. Our goal was simply to spell out that the speculations on their possible breakdown for the classical Heisenberg model, or for the one-parameter scaling theory of localization, are based on an unreasonable mathematical analysis.

We acknowledge the support by CNRS-JSPS cooperative research project, and by CREST of Japan Science and Technology Corporation.

References

- [1] A. M. Polyakov, Phys. Lett. **59 B**, 79 (1975).
- [2] E. Brézin and J. Zinn-Justin, Phys. Rev. Lett. **36**, 691 (1976).
- [3] K. G. Wilson and M. E. Fisher, Phys. Rev. Lett. **28**, 240 (1972).
- [4] P. W. Anderson, Phys. Rev. **109**, 1492 (1958).
- [5] F. Wegner, Z. Phys. **B 35**, 207 (1979).
- [6] E. Brézin, S. Hikami and J. Zinn-Justin, Nucl. Phys. **B 165**, 528 (1980).
- [7] S. Hikami, Phys. Rev. **B 24**, 2671 (1981).
- [8] V. E. Kravtsov, I. V. Lerner and V. I. Yudson, Sov-Phys. JETP **67**, 1441 (1988).
- [9] F. Wegner, Z. Phys. **B 78**, 36 (1990).
- [10] K. Wilson and J. Kogut, Phys. Report **12C**, 75 (1974).
- [11] G. E. Castilla and S. Chakravarty, *Is the phase transition in the Heisenberg model described by the $2 + \epsilon$ expansion of the non-linear σ model?* preprint, cond-mat/9605088.
- [12] A. N. Vasil'ev and A. S. Stepanenko, Theor. Math. Phys. **95**, 471 (1993).
- [13] S. K. Ma, Phys. Rev. **A 10**, 1818 (1974).

- [14] S. Hikami, Prog. Theor. Phys. **64**, 1466 (1980).
- [15] E. Abrahams, P. W. Anderson, D. C. Licciardello and T. V. Ramakrishnan, Phys. Rev. Lett. **42**, 673 (1979).
- [16] E. Brézin and S. Hikami, in preparation.